

Confocal Microscopic Analysis of Integrin Expression on the Microvasculature and its Sprouts in the Neonatal Foreskin

Judy Enenstein* and Randall H. Kramer*†

*Departments of Anatomy and Stomatology and the †Cardiovascular Research Institute, University of California San Francisco, San Francisco, California, U.S.A.

Members of the integrin family of adhesion receptors are essential participants in blood vessel growth and remodeling. It is not known which integrins are involved in the initial stages of angiogenesis *in vivo*. In this study we determined the location of integrins on the blood vessels of a growing tissue, the neonatal foreskin, in which neovascularization is likely to occur. We used the confocal microscope to visually reconstruct vessels from the papillary dermis of the foreskin and to identify potential sprouts as narrow, tapering extensions from these vessels. Blood vessels were initially identified by their positive reaction with antibodies to von Willebrand factor or human platelet endothelial cell adhesion molecule and their negative response to anti-neurofilament antibodies. Later, vessels were identified by their shape and location. We screened

vessels with anti bodies to integrin subunits $\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 5$, $\alpha 6$, αv , $\beta 1$, $\beta 3$ and $\beta 4$. We found that integrin subunits $\alpha 6$ and $\beta 4$ were consistently found along the whole length of capillary loops and extended to the distal ends of presumed sprouts. The $\alpha 2$ and αv integrin concentrations, which are normally low in the microvasculature, were increased on the sprouts. $\alpha 5$ was either absent from vessels entirely or more concentrated on the body than on the sprout. $\alpha 1$ was more commonly present on nerves than blood vessels. These studies suggest an important role for the $\alpha 6\beta 4$ integrin in the initial stages of endothelial outmigration during new vessel growth. **Key words:** endothelial sprout/adhesion/angiogenesis/extracellular matrix. *J Invest Dermatol* 103:381–386, 1994

Angiogenesis, whether during development or wound healing, entails the sprouting of new vessels from pre-existing vessels. Endothelial cells extend processes through breaks in the basement membrane and then migrate as a column into the interstitial matrix. This migration consists of pairs of endothelial cells sliding past each other. Proximally the column may contain a lumen, but distally it is a solid cord [1–3].

Cell migration is a complex process requiring adhesion receptors to provide traction without binding cells too tightly to the substratum, and intracellular signaling to realign the cytoskeleton and initiate cortical flow [4]. Members of the integrin family of heterodimeric transmembrane receptors are involved in cell-substratum and cell-cell adhesion and also initiate various intracellular signals. The integrins bind to extracellular matrix molecules through their external domain, and may directly attach to cytoskeletal elements (talin and α -actinin) via their cytoplasmic domain. The ligand specificity and signaling ability of integrins are determined by varying the combination of α and β subunits that make up the heterodimers. There are twelve known α subunits and eight known β subunits. The integrin subunits $\alpha 1$, $\alpha 2$, $\alpha 3$, $\alpha 5$, $\alpha 6$, $\beta 1$, $\beta 3$, $\beta 4$, and $\beta 5$ have been detected on endothelial cells [5–9].

Angiogenesis obliges endothelial cells to leave the collagen IV/laminin-rich basement membrane and migrate through a matrix rich in interstitial collagen and fibronectin. We assumed that this change of substratum would require a different array of integrins on the endothelial sprout. Our purpose was to determine which integrins are important in the initial stages of blood vessel growth and repair. To do so, we have used confocal microscopy to reconstruct vessels from laser sections and identify sprouts as thin cords arising from vessels.

The confocal microscope, which allows the use of thick tissue sections (10 μ or more are visually sectioned down to 0.5–1.0 μ and then reconstructed by computer), has several advantageous features for examining sprouts. Because vessels can be studied over longer distances than usual, a) sprouts can be visualized, b) molecules can be compared by double-labeling immunofluorescence for their location along the sprout, and c) chances of “sprout” artifacts are greatly decreased because serial reconstruction shows whether a thin strand is truly a vessel outgrowth or only part of a longer vessel. In addition, the thick sections diminish the possibility that a vessel “end” will be created by cutting off the rest of the vessel.

We used double-labeling immunofluorescence microscopy to compare the location of various integrins on vessel sprouts with the location of the $\alpha 6$ integrin. The $\alpha 6$ integrin was chosen as a standard because antibodies to $\alpha 6$ produce intense staining of all endothelial cells at blood vessel basement membranes [9], and because we noted that this staining appeared to extend far into sprouts. The $\alpha 6$ integrin is also found on the epidermis and glands, and on infiltrating cells such as macrophages, but is not found on fibroblasts or smooth muscle cells by tissue staining [10]. Our studies indicate that integrins $\alpha 6$ and $\beta 4$ are found on the narrow extensions of sprouts

Manuscript received August 19, 1993; accepted for publication April 6, 1994.

Reprint requests to: R. Kramer, University of California, Room HSW-604, Box 0512, San Francisco, CA 94143-0512.

Abbreviation: PECAM, human platelet endothelial cell adhesion molecule.




Figure 1. Hematoxylin and eosin stain of neonatal foreskin. Arrow indicates capillary tip approaching epidermis (E).

and may show a concentration at the distal tip. $\alpha 2$ and αv , though less concentrated than $\alpha 6$, are also found along the sprouts. Integrins $\alpha 5$ and $\beta 3$ are more prevalent on the body of blood vessels, and are entirely absent from some vessels.

MATERIALS AND METHODS

Tissue Preparation Human foreskin was obtained from the Well-baby Nursery at the University of California San Francisco. Tissues were fixed for 4 h in 4% formaldehyde in phosphate-buffered saline (PBS), transferred to 15% sucrose for 1 h, embedded in O.C.T. (Miles Laboratories, Elkhart Lake, Indiana), and frozen in liquid nitrogen. Tissue blocks were stored at -80°C . Frozen sections 10–18 μ thick were placed on superfrost glass slides (Fisher).

Immunofluorescent Staining All incubations and washes were done at room temperature in 1% goat serum in PBS. Nonspecific staining in tissue sections was blocked by incubating for 1 h in 1% goat serum in PBS. Sections were then treated with primary antibodies for 20 min, rinsed five times, treated with secondary antibodies conjugated to fluorescein (at 1:50) or rhodamine-lissamine (at 1:1000) for 20 min, washed five times, and mounted in Fluoromount-G (Fisher Biotech, Pittsburgh, PA). In all cases shown, $\alpha 6$ was labeled with rhodamine.




Figure 2. Expression of $\alpha 6$ on PECAM-positive vessels. Projection of confocal Z series. Neonatal foreskin was simultaneously labeled by indirect immunofluorescence with antibodies to a) the endothelial cell marker PECAM (with fluorescein secondary) and b) the $\alpha 6$ integrin (with rhodamine secondary). $\alpha 6$ is found both on blood vessels and on the dermal-epidermal junction (J). Note narrowing of vessel near tip (arrows). E, epidermis. Bar, 10 μm .




Figure 3. Neurofilament localization in foreskin. Projection of confocal Z series. A) The shape of fluorescein-labeled neurofilaments (a) is distinct from that of rhodamine-labeled vWF-positive blood vessels (b). Arrowheads, nerves. Hollow arrows, blood vessels. B) Fluorescein-labeled neurofilaments (a) are distinct from rhodamine-labeled $\alpha 6$ extensions (b). Arrow, tip of vessel extension. E, epidermis. J, dermal-epidermal junction. Bar, 10 μm .

Antibodies Mouse anti-human $\alpha 1$ was from T-Cell Sciences, Cambridge, MA. Mouse monoclonal anti- $\alpha 2$ (12F1) was provided by Dr. Virgil Wood, University of California, San Diego, CA [11]. Mouse monoclonal anti- $\alpha 3$ (VM-2) was provided by Dr. Vera Morhenn, SRI International, Menlo Park, CA [12]. Rabbit antibody to $\alpha 5$ was provided by R. Hynes, MIT, Boston [13]. Rat monoclonal anti- $\alpha 6$ (J1B5) was provided by Dr. Caroline Damsky, UCSF, San Francisco, CA [14]. Mouse monoclonal anti-human integrin αv subunit (LM142) was provided by Dr. David Cheresh, Research Institute of Scripps Clinic, La Jolla, CA [15]. Mouse monoclonal anti-human integrin $\beta 1$ (A-1A5) was provided by Dr. Martin Hemler, Dana Farber Cancer Institute, Boston, MA [16]. Rabbit antibody to human β (anti-GpIIb-IIIa) was provided by Dr. David Phillips, UCSF [17]. Mouse monoclonal anti- $\beta 4$ (3E1) and polyclonal anti-human laminin were purchased from Telios Pharmaceuticals, San Diego, CA. Mouse monoclonal anti-epiligrin (GB3) was purchased from Accurate Chemical and Scientific Corporation, Westbury, NY. Polyclonal antibodies to von Willebrand factor (vWF) were purchased from Sigma Chemical, St. Louis, MO. Monoclonal antibodies to neurofilaments (200 kDa protein) were from Boehringer Mannheim, Indianapolis, IN. Monoclonal antibodies to human platelet endothelial cell adhesion molecule (PECAM) were from R and D Systems, Minneapolis, MN. Goat anti-mouse, anti-rat and anti-rabbit IgG conjugated with fluorescein-isothiocyanate and goat anti-mouse, anti-rat, and anti-rabbit IgG conjugated with lissamine-rhodamine were from Jackson Immuno-Research, West Grove, PA. Antibodies were tested to ensure that there was

no crossreactivity of anti-mouse secondary antibody with rat primary antibody or vice versa.

Microscopy Samples were viewed on a Biorad MRC 600 confocal microscope and analyzed with the COMOS program. The papillary dermis was searched for potential vessel sprouts (areas of vessel narrowing). These areas were visually sectioned to 1 μ by the microscope. The dermal-epidermal junction was used as a marker for determining tissue thickness. Sections were reconstructed into two-dimensional projections from a 1- μ series. All micrographs are projections of a Z series unless otherwise indicated.

RESULTS

We chose human neonatal foreskin in which to study microvascular development because there is significant neovascularization in skin during the first three postnatal months [18–20], and because a large number of probes are available for human integrins and their ligands. After birth, the disorganized subpapillary plexus in the skin is remodeled into discrete pathways, and capillary loops are projected deep into the dermal papilla. **Figure 1** shows a hematoxylin-and-eosin-stained section of neonatal foreskin. Note how the capillary tapers as it extends towards the epidermis.

We used both immunologic and morphologic criteria to identify sprouts arising from the microvasculature. First, blood vessels were identified within the dermis of human foreskin by their reaction with antibodies to the endothelial cell markers vWF or PECAM. Sprouts were recognized as tapering continuous extensions from these vessels that were verified not to be fragmented vessels by confocal sectioning. Whereas capillaries are typically 5–10 μ m in diameter, the sprout tips are solid cords that narrow to ~2 μ m or sometimes have wider club-like shapes [2]. The confocal microscope allows the observer to follow vessels through a thick tissue block and to distinguish the top and bottom of sections of the block. We consider that those extensions that taper and end in the middle sections of a series are likely to be sprouts and not cutting artifacts.

Our initial studies compared the location of various integrins with that of the blood vessel markers PECAM and vWF. **Figure 2** is a confocal two-dimensional reconstruction of neonatal foreskin double-labeled for PECAM and α 6 by indirect immunofluorescence. The blood vessel tapers until it is too narrow to contain a lumen. PECAM colocalizes with α 6 on both the vessel and the tapering extension. However, the dermal-epidermal junction is

α 6-positive and PECAM-negative. Comparison of other integrins with PECAM or vWF confirmed previous reports [5,9,21] that α 2, α 3, α 5, α v, β 1 and β 4 are present on the body of some blood vessels of the microvasculature (not shown).

Because α 6 stains blood vessels continuously and intensely, we chose α 6 as the standard with which to compare the location of other integrins. However, because α 6 is also found on nerves, we first compared the location and appearance of nerves, as labeled by antibodies to the 200-kDa neurofilament protein, with that of vWF or α 6 in the papillary dermis (**Fig 3**). Unlike the blood vessels, which taper into sprouts, the nerves remain very narrow over long distances. Sometimes the nerve fiber twists around a vessel or runs alongside a blood vessel. **Figure 3B** shows a tapering α 6-positive vessel that does not react with anti-neurofilament antibodies.

Comparison of other integrins with α 6 revealed that β 4 colocalized on sprouts with α 6 (**Fig 4**). **Figure 4B** shows part of a Z series of the same tissue block. Note that the narrow vessel extension is longest in section 5 and gradually diminishes in length in subsequent sections. Like α 6, antibodies to β 4 stained basement membranes of endothelium and epithelium intensely. Whereas α 6 and β 4 were present along most of the vessel length, staining for both α 2 and α v was pale on much of the vessel body, but was sometimes concentrated in the distal part of the vessel containing the sprout (**Fig 5A,B**). The diffuse staining of α 2 and α v suggested cytoplasmic localization rather than the basement membrane localization of α 6. Antibody to α 2 usually stained the basal epidermal cells much more intensely than the blood vessels (**Fig 5A**) whereas α v staining was patchy or faint in the epidermis (**Fig 5B**). Antibody to β 1, like that to α 6, stained the full length of many vessels, but the two stains did not always colocalize either along the vessel or at the tip (**Fig 5C**). Areas in which β 1 and α 6 did not colocalize likely represented α 6 β 4-rich sites. It was also noted that staining of the epidermis with anti- β 1 antibodies produced a circumcellular pattern, without a distinct basement membrane stain, suggesting a lack of α 6 β 1 on the basement membrane. Anti- β 3 stained some vessel bodies brightly (**Fig 5D**) and others not at all; it also extended along some sprouts (not shown). Three integrin subunits were almost never seen on vessel sprouts. Staining for the α 5 integrin was either more concentrated on the vessel body than on tapering extensions or missing entirely (**Fig 5E**). α 5 was only infrequently seen on epidermal patches. Staining for α 3, which was pale on blood vessels, was




Figure 4. Colocalization of β 4 and α 6 in projection of Z series and individual Z-series sections. **A**) Projection of Z series. Fluorescein-labeled β 4 (**a**) colocalizes with rhodamine-labeled α 6 (**b**). Arrow, tip of vessel extension. E, epidermis. J, dermal-epidermal junction. **B**) 1 μ sections from the Z series. Sections 2–10 of a 12-section series are shown. Section 5 corresponds to the longest tapering extensions shown in **A**. Bar, 10 μ m.




Figure 5. Localization of $\alpha 2$, $\alpha 5$, $\beta 1$, $\beta 3$, $\alpha 5$, and $\alpha 1$ on blood vessels. Projection of confocal Z series. Foreskin was double-labeled with fluorescein for either $\alpha 2$, $\alpha 5$, $\beta 1$, $\beta 3$, $\alpha 5$, or $\alpha 1$ and with rhodamine for $\alpha 6$. **A)** $\alpha 2$ is prominent in basal epidermal cells and diffusely present on the end of blood vessel extensions (a) as compared with $\alpha 6$ (b). **B)** $\alpha 5$ is diffusely present near the vessel tip (a) as compared with $\alpha 6$ (b). Note the break in staining just before the tip on the $\alpha 6$ -positive sprout. This break is characteristic of $\alpha 6$ in sprouts. **C)** $\beta 1$ is brightly stained along the vessel length (a) but is not always colocalized with $\alpha 6$ (b). **D)** $\beta 3$ is present on the blood vessel body (a) lesser amounts on the vessel extension, visualized by staining $\alpha 6$ (b). **E)** $\alpha 5$ is sometimes found along blood vessels (a) but is less distinct on vessel extensions, as visualized by staining $\alpha 6$ (b). **F)** $\alpha 1$ is predominantly found on the nerve (a) encircling the $\alpha 6$ -positive vessel (b). Epidermis in F is on the right side of picture. E, epidermis. J, dermal-epidermal junction. Arrows, tips of vessel extensions. Bar, 10 μ m.

concentrated on basal epidermal cells (not shown). $\alpha 1$ was more prevalent on nerves than on blood vessels (Fig 5F).

We also compared the localization of $\alpha 6$ with that of its potential ligands: laminin, and the laminin variant epiligrin. Epiligrin was usually not present on blood vessels. A-chain laminin was present on all $\alpha 6$ -positive vessels, as were its individual A and B subunits (data not shown).

DISCUSSION

Using the confocal microscope to reconstruct blood vessels, we have examined the localization of various integrins along the vessels and their newly forming sprouts. The integrins $\alpha 6$ and $\beta 4$ were most consistently found on both the body of microvascular vessels and on the long, tapering sprouts. Integrins $\alpha 2$ and αv sometimes colocalized with the tips of these extensions, whereas $\alpha 5$, when present, was denser on the vessel body than on the distal end of the sprout.

The known ligands for $\alpha 6\beta 1$ and $\alpha 6\beta 4$ are members of the laminin family. The consistent localization of $\alpha 6$ and $\beta 4$ at the distal ends of sprouts came as a surprise, because endothelial cells leave the laminin/type IV collagen-rich blood vessel basement membrane during angiogenesis and migrate through a matrix rich in fibronectin, fibrin, and interstitial collagen. However, studies of induced angiogenesis in corneas have detected laminin at the tips of newly forming vessels, and even on individual cells in advance of the tip [22,23]. When human microvascular endothelial cells are plated on fibronectin *in vitro*, they synthesize laminin within 24 h [23]. In addition, the endothelial cell line EAHY 926 forms tubes in the laminin-rich matrix Matrigel. Significantly, this tube formation is blocked by antibodies to $\alpha 6$ or $\beta 1$, but not by antibodies to $\alpha 5$. Blockage occurs when as few as 10% of $\alpha 6$ or $\beta 1$ sites are bound by antibody [24]. Much higher antibody concentrations are needed to block cell adhesion. Drake *et al* [25] provided evidence that an earlier stage of angiogenesis, cord formation, is not dependent on $\beta 1$ integrins. They showed that when chick embryos are treated with the CSAT antibody, which perturbs $\beta 1$ integrins, formation of blood vessel tubes from solid cords is blocked, but initial cord formation is unaffected. Therefore, if integrins function in cord formation *in vivo*, one would expect that a major component would be some non- $\beta 1$ integrin, such as $\beta 4$.

$\alpha 6\beta 4$ is primarily associated with epiligrin in hemidesmosome complexes on epithelial cells [26]. However, well-developed hemidesmosomes are not present in the microvasculature. Because epiligrin (A. Sonnenberg, personal communication) and A-chain laminin [27] are ligands of $\alpha 6\beta 4$ in other cell types, we attempted to stain for epiligrin with GB3 antibodies. Results were usually negative. Either epiligrin is not present or microvascular epiligrin is not accessible to the GB3 antibodies. A third possibility is that a variant epiligrin-like laminin isoform exists that could bind $\alpha 6\beta 4$. In fact, a truncated variant of laminin has been detected in aortic endothelial cells [28]. We have observed that dermal microvascular endothelial cells do not make significant A-chain laminin, but seem to synthesize a truncated form of laminin (unpublished observations).

Although $\alpha 6$ and $\beta 4$ were present along the whole length of the vessel and sprout, $\alpha 2$ and αv staining was relatively weak along the vessel body and was sometimes detectable only on the distal end of the vessel, including the sprout. The increased density of $\alpha 2$ and αv on vessel sprouts is evidence that they are active in angiogenesis. Previous *in vivo* studies have shown that both $\alpha 2$ and αv are present during capillary growth. $\alpha 2$ levels are high during embryonic glomerular capillary formation but low in capillaries in adults [29], and $\alpha v\beta 3$ has been detected in human granulation tissue [21]. Concentrations of $\alpha 2$ and αv , as well as $\alpha 5$, are upregulated by the angiogenic factor basic fibroblast growth factor (bFGF) [30,31]. *In vitro* tube formation studies provide complex information on the role of $\alpha 2$ and αv . Human umbilical vein endothelial cells (HUVEC) plated on Matrigel were inhibited from forming tubes in the presence of anti- αv antibodies [32], whereas HUVEC plated on fibrin were stimulated to produce more and longer tubes in the presence of antibodies to $\alpha v\beta 3$ and phorbol myristic acid [33]. Likewise,

HUVEC plated on collagen were stimulated by antibodies to $\alpha 2$. However, αv and $\alpha 2$ antibodies were not stimulatory when the substrates were reversed (HUVEC were plated on collagen and fibrin, respectively). Thus, *in vitro* tube studies cannot yet predict which integrins are important during *in vivo* angiogenesis, a process that depends on a complex and changing mixture of ligands.

The finding that $\alpha 5$ and $\beta 3$ are present only on a subset of microvascular endothelia is consistent with previous studies in which $\alpha 5\beta 1$ and $\alpha v\beta 3$ were present in granulation tissue but not quiescent blood vessels [21]. This in turn provides further evidence that the microvasculature in the neonatal foreskin is undergoing active remodeling. The presence of the fibronectin receptor, $\alpha 5\beta 1$, on wider parts of vessels but not the narrowest extensions suggests that it may be important at a later stage of vessel development, perhaps as cord cells spread and form tubes. Nicosia *et al* [34] found that when rat aortas were embedded in collagen gels, the length of outgrowing microvessels increased with increasing doses of fibronectin, but the number of vessels did not increase. This implies that fibronectin does not stimulate the initial outgrowth of new vessels.

We used several criteria to identify blood vessel sprouts. Sprouts were initially recognized as tapering extensions from PECAM- or vWF-positive vessels. To screen out cutting artifacts, we considered only those extensions that arose in the middle of the confocal section series as sprouts. Because tissues were fixed before sectioning, it is unlikely that membranes vesiculated or solubilized. Nerves were initially identified by their reaction with antibodies to the 200-kDa neurofilament protein and subsequently by their shape and location. Nerves in the papillary dermis are generally thinner than blood vessels and extend for long distances without tapering. Because nerves may also wrap around blood vessels, we carefully examined confocal sections to ensure that the "sprout" extensions arose continuously from the full body of the vessel, rather than from a thin fiber running alongside. Extensions from distinctly bipolar or multiarmed cells were likewise discounted as potential sprouts, because they were likely to be pericytes or dermal cells. Remodeling of the neonatal circulation entails both growth and regression of blood vessels. Without a marker for regression, it is impossible to be certain whether a vessel is growing or regressing. Despite careful elimination of alternative possibilities, we recognize that what we present is the strong suggestion of blood vessel sprouts. Without a definitive chemical marker of sprouts, proof of the location of various integrins on vessel sprouts awaits an immunoelectronmicroscopic study.

What we have described is a static vision of the vessel sprouts. Although $\alpha 6$ and $\beta 4$ were consistently the most intensely staining integrins at the ends of sprouts, this does not diminish the importance of other integrins, which may be present in low concentration and/or appear transiently at critical moments of angiogenesis. The integrin concentration needed to respond to the original angiogenic stimuli, and perhaps to recruit other integrins, may be much smaller than that needed to anchor the sprout physically. Whether $\alpha 6$ and $\beta 4$ provide both signals and adhesiveness has yet to be determined.

Note Added in Proof: Recently it has been reported that $\alpha v\beta 3$ integrin is required for angiogenesis in the chick chorioallantoic membrane (Brooks PC, Clark RA, Cheresh DA: Requirement of vascular integrin $\alpha v\beta 3$ for angiogenesis. *Science* 264:569–571, 1994).

We thank Evangeline Leash for her invaluable editorial assistance. We thank Dr. Ron Clyman for very helpful comments on the manuscript. This work was supported by a research fellowship from the American Heart Association, California Affiliate, 91-08, with funds contributed by the Alameda County Chapter; by AHA grant 880763; and by NIH grants CA33834, CA51884, and DE-00242.

REFERENCES

1. Folkman J, Klagsbrun M: Angiogenic factors. *Science* 235:442–447, 1987
2. Rhodin JAG, Fujita H: Capillary growth in the mesentery of young rats. Intravital video and electron microscope analysis. *J Submicrosc Cytol Pathol* 21:1–34, 1989

3. Paku S, Paweletz N: First steps of tumor-related angiogenesis. *Lab Invest* 65:334-346, 1991
4. Stossel TP: On the crawling of animal cells. *Science* 260:1086-1094, 1993
5. Albelda SM, Daise M, Levine EM, Buck C: Identification and characterization of cell-substratum adhesion receptors on cultured endothelial cells. *J Clin Invest* 83:1992-2002, 1989
6. Cheng Y-F, Kramer R: Human microvascular endothelial cells express integrin-related complexes that mediate adhesion to the extracellular matrix. *J Cell Physiol* 139:275-286, 1989
7. Kennel SJ, Vodfrey V, Ch'ang LY, Lankford TK, Foote LJ, Maddinje, A: The $\beta 4$ subunit of the integrin family is displayed on a restricted subset of endothelium in mice. *J Cell Science* 101:145-150, 1992
8. Pasqualini R, Bodorova J, Ye S, Hemler ME: A study of the structure, function and distribution of $\beta 5$ integrins using novel anti- $\beta 5$ monoclonal antibodies. *J Cell Science* 105:101-111, 1993
9. Ryyanen J, Jaakkola S, Engvall E, Peltonen J, Uitto J: Expression of $\beta 4$ integrins in human skin: comparison of epidermal distribution with $\beta 1$ -integrin epitopes, and modulation by calcium and vitamin D3 in cultured keratinocytes. *J Invest Dermatol* 97:562-567, 1991
10. Sonnenberg A, Linders CJT, Daams JH, Kennel SJ: The $\alpha 6 \beta 1$ (VLA-6) and $\alpha 6 \beta 4$ protein complexes: tissue distribution and biochemical properties. *J Cell Sci* 96:207-217, 1990
11. Pischel KD, Hemler ME, Huang C, Bluestein HG, Woods VL: Use of the monoclonal antibody 12F1 to characterize the differentiation antigen VLA-2. *J Immunol* 138:226-233, 1987
12. Morhenn VB, Roth S, Roth R: Use of a monoclonal antibody (VM-2) plus the immunogold-silver technique to stain basal cell carcinoma cells. *J Am Acad Dermatol* 17:765-769, 1987
13. Hynes RO, Marcantonio EE, Stepp MA, Urry LA, Yee GH: Integrin heterodimer and receptor complexity in avian and mammalian cells. *J Cell Biol* 109:409-420, 1989
14. Werb Z, Tremble PM, Behrendtsen O, Crowley E, Damsky CH: Signal transduction through the fibronectin receptor induces collagenase and stromelysin gene expression. *J Cell Biol* 109:877-889, 1989
15. Cheresch DA, Spiro RC: Biosynthetic and functional properties of an Arg-Gly-Asp directed receptor involved in human melanoma cell attachment to vitronectin, fibronectin, and von Willebrand factor. *J Biol Chem* 262:17703-17711, 1987
16. Hemler ME, Ware CF, Strominger JL: Characterization of a novel differentiation antigen complex recognized by a monoclonal antibody (A-1A5): unique activation-specific molecular forms on stimulated T cells. *J Immunol* 131:334-340, 1983
17. Charo IF, Bekeart LS, Phillips DR: Platelet glycoprotein IIb-IIIa-like proteins mediate endothelial cell attachment to adhesive proteins and the extracellular matrix. *J Biol Chem* 262:9935-9938, 1989
18. Tonnesen MG, Jenkins D, Siegal SL, Lee LA, Huff JC, Clark RAF: Expression of fibronectin, laminin, and factor VIII-related antigen during development of the human cutaneous microvasculature. *J Invest Dermatol* 85:564-568, 1985
19. Perara P, Kurban AK, Ryan TJ: The development of the cutaneous microvascular system in the newborn. *Br J Dermatol* 82(suppl 5):86-91, 1970
20. Wolff JR, Goerz C, Bar T, Guldner FH: Common morphogenetic aspects of various organotypic microvascular patterns. *Microvasc Res* 10:373-395, 1975
21. Rudolf R, Cheresch D: Cell adhesion mechanisms and their potential impact on wound healing and tumor control. *Clin in Plast Surg* 17:457-462, 1990
22. Form DM, Pratt BM, Madri JA: Endothelial cell proliferation during angiogenesis. In vitro modulation by basement membrane components. *Lab Invest* 55:521-530, 1986
23. Jerdan JA, Michels RG, Glaser BM: Extracellular matrix of newly forming vessels—an immunohistochemical study. *Microvasc Res* 42:255-265, 1991
24. Bauer J, Margolis M, Schreiner C, Edgell C-J, Azizkan J, Lazarowski E, Juliano RL: In vitro model of angiogenesis using a human endothelium-derived permanent cell line: contributions of induced gene expression, G-proteins and integrins. *J Cell Physiol* 153:437-449, 1992
25. Drake CJ, Davis LA, Little CD: Antibodies to $\beta 1$ integrins cause alterations of aortic vasculogenesis, in vivo. *Dev Dynamics* 193:83-91, 1992
26. Carter WG, Ryan MC, Gahr PJ: Epiligrin, a new cell adhesion ligand for integrin $\alpha 3 \beta 1$ in epithelial basement membranes. *Cell* 65:599-610, 1991
27. Lee EC, Lotx MM, Steele GD, Mercurio AM: The integrin $\alpha 6 \beta 4$ is a laminin receptor. *J Cell Biol* 117:671-678, 1992
28. Tokida Y, Aratani YY, Morita A, Kitagawa Y: Production of two variant laminin forms by endothelial cells and shift of their relative levels by angiostatic steroids. *J Biol Chem* 265:18123-18129, 1990
29. Korhonen M, Ylanne J, Laitinen L, Virtanen I: The $\alpha 1$ - $\alpha 6$ subunits of integrins are characteristically expressed in distinct segments of developing and adult human nephron. *J Cell Biol* 111:1245-1254, 1991
30. Enenstein J, Walch NS, Kramer RH: Basic FGF and TGF- β differentially modulate integrin expression of human microvascular endothelial cells. *Exp Cell Res* 203:499-503, 1992
31. Swerlich RA, Brown EJ, Xu U, Lee DH, Mianos S, Lawley TJ: Expression and modulation of the vitronectin receptor on human dermal microvascular endothelial cells. *J Invest Dermatol* 99:715-722, 1992
32. Davis CM, Daneshmandi SC, Laurenza A, Molony JL: Identification of a role of the vitronectin receptor and protein kinase C in the induction of endothelial cell vascular formation. *J Cell Biochem* 51:206-218, 1993
33. Gamble JR, Matthia LJ, Meyer G, Kaur P, Russ G, Faull R, Berndt MC, Vadas MA: Regulation of in vitro capillary tube formation by anti-integrin antibodies. *J Cell Biol* 121:931-943, 1993
34. Nicosia RF, Bonanno E, Smith M: Fibronectin promotes the elongation of microvessels during angiogenesis in vitro. *J Cell Physiol* 154:654-661, 1993